

PREVALENCE AND ASSOCIATED FACTORS OF MEDICATION NON-ADHERENCE IN HEMATOLOGICAL-ONCOLOGICAL PATIENTS IN THEIR HOME SITUATION

Linda Bouwman,
Corien M. Eeltink, Otto Visser,
Jeroen J. W. M. Janssen and
Jolanda M. Maaskant

DISCLOSURE PRESENTER




- ✘ No (potential) conflict of interests
- ✘ 1. Relations that could be relevant for the meeting
- ✘ 2. Sponsorship or research funds
- ✘ 3. Payment or other (financial) remuneration
- ✘ 4. Shareholder
- ✘ 5. Other relation

CONCLUSION

- Prevalence 50%
- Associated factors
 - Fatigue
 - Higher education level
 - Younger age

NON-ADHERENCE

A deviation from the prescribed medication regimen sufficient to adversely influence the regimen's intended effect



METHOD

MARS-5

	Always	Often	Sometimes	Rarely	Never
I decide to miss out a dose					
I forget to take it					
I alter a dose					
I stop taking it for a while					
I take less than instructed					

BMQ, HADS, ICQ, EORTC-QLQ-30

socio-economic factors and disease

RESULTS

- 472 questionnaires were sent
- 280 returned
- 259 included

CONCLUSION

- Prevalence 50%
- Associated factors
 - Fatigue
 - Higher education level
 - Younger age

QUESTIONS



Lbouwman@sparnegasthuis.nl